YDEF training survey 2014

Paul Grant, Ali Chakera, Kelly Cheer, Roselle Herring, Rahat Maitland, Jyothis George, Milan Piya, Stuart Little

YOUNG DIABETOLOGISTS & ENDOCRINOLOGISTS

Introduction

The remit of the YDEF – an organised collective of registrars and newly appointed Consultants from across the UK – is to promote 'Education, Representation, and Communication'. In addition to providing educational courses and events, YDEF seeks to improve the quality of specialist education by identifying deficiencies in training, and reporting these to those responsible for organising that training, both locally and nationally, with constructive suggestions for improvements.¹

Key concerns in the past have been access to training opportunities in elements of the D&E curriculum such as insulin pump experience or specialist pituitary clinics,² marked variation in the quality and types of training between centres and the different regions, and the constant dilution of specialist training opportunities by the increasing service demands of general internal medicine – the management of the acute take, and of general medical inpatients.

D&E specialist trainees typically undertake training in GIM in parallel with D&E leading to dual certification. This is not the model for all medical specialties, so the available pool of registrars expecting to undertake general medical work (and training) has contracted, and often comprises only D&E and Elderly Medicine trainees. It has been widely acknowledged that the role of a medical registrar has become increasingly challenging and potentially dispiriting.3 As a result of national targets and financial pressures, hospitals are forced to prioritise acute 'general medical' issues over long-term outpatient based specialist care. Coupled to this is the fact that there have been significant changes in the speciality over the years (i.e. increased insulin pump use, expansion of community diabetes care and the introduction of a Speciality Certificate Examination). In addition training schemes are struggling to fill vacancies, which further aggravates the situation, impacting negatively on the perception of D&E as a career. These issues have been acknowledged nationally and addressed in recent reports.4,5

There has not been a comprehensive national training survey for some time^{6,7} and it is important to obtain an up to date 'state of the nation' picture of current issues to

ensure that the YDEF's activities, as well as those of other bodies responsible for providing and improving specialist D&E training, remain focused on the needs of trainees.8

Methods

The YDEF Training Survey was launched at its annual meeting in March 2014. It was modelled on the recent oncology registrars' training study.⁹ It included questions covering domains of training such as the balance of time spent between different specialities, the demands of service provision, experience of appraisal and assessment, availability and uptake of study leave, time spent 'out of programme' and measures of the quality and satisfaction with practical experience in the workplace. In addition to quantitative data, respondents were encouraged to add free text responses to explain their answers and provide qualitative information.

Results

There are currently 463 doctors in accredited D&E higher specialist training posts in the UK; 94 (20%) completed the survey.

 Balance of training: trainees reported that at most a third of their time was spent in D&E (Figure 1) with 68% of their time taken up by GIM. Specific comments from trainees relating to this perceived lack of balance in their training included: 'Heavy commitment towards GIM', 'Ser-


Abbreviations and acronyms

CCT certificate of completion of training
D&E diabetes and endocrinology
DAFNE Dose Adjustment for Normal Eating
GIM general internal medicine
MDT multidisciplinary team
YDEF Young Diabetologists &
Endocrinologists' Forum

vice provision is taking over' and 'Bias to GIM – it seems like we get a lot less training time than other specialities'.

- Clear training plan: when starting in a new training post a clear training plan had been provided to 76% of respondents and 65% had received an appraisal at the beginning or end of their clinical attachment. The majority (81%) had found this a constructive process.
- Study leave and budget: trainees are allocated 30 days study leave per year, but only 72% of trainees were able to take the study leave they needed. The majority spent all of their study budget (approx. £800), spending 60% and 40% on D&E and GIM respectively.
- Training days: registrars were happy (77%) with access to training days but noted the variability in quality of content

 the average training day rating was only 6.6 out of 10, with the major criticism being an excess of 'filler' content.



- When asked 'What are the main deficiencies in your training?' Responses were grouped into similar themes outlined in Table 1; the majority of respondents (55-69%) reported that they were able to discuss a range of issues with their educational or training programme supervisors and that they were able to configure their training rotation to suit their requirements.
- Ouality of clinical experience: trainees ranked individual aspects of their D&E training exposure in order to gauge not the volume of exposure but the depth and the quality. Results varied, with outpatient endocrinology experience faring better than diabetes outpatients (54% good or excellent vs only 33% good or excellent for diabetes). Fewer respondents thought their diabetes inpatient experience was good (20% good or excellent vs 25% poor or very poor ratings), however 71% of respondents were more than satisfied with the quality of their MDT experiences. Figure 2 provides a detailed breakdown of how respondents viewed different training areas. Only a minority (18%) of respondents were dissatisfied with their training overall (Figure 3).
- In response to the question, "What do you consider the best aspects of your training?", some of the free-text comments included; 'having time to enjoy and understand my speciality', weekly case discussions', 'free courses from the YDEF', 'academic opportunities', endocrine clinics and post clinic meetings', 'courses and conferences' and 'interesting case mix'.
- Engagement in audit / research / service improvement: clinical research was generally considered an extracurricular activity, but 24% of responders had experience of clinical research, even fewer (14%) had been involved in service improvement (e.g. introducing a diabetes admissions avoidance plan¹⁰ or an insulin safety group¹¹), but 82% had undertaken obligatory audits. It is noteworthy that several trainees (49%) were pursuing additional qualifications (Figure 4). Reasons for this included: academic interest, personal interests/ curiosity, opportunity for career promotion, opportunities to have children and to improve one's job prospects/CV.


Discussion

Questionnaire surveys of doctors in training often suffer from poor response rates, marked

Table 1 Trainee comments relating to perceived problems with training


Theme	Examples
Lack of time	'priority is service provision' 'need more time in D&E'
Lack of structured teaching	'educational supervisors need more insight and training' 'no clear structure' 'lack of support and mentorship'
Sub-speciality exposure	'minimal insulin pump access' 'lack of paediatric and transition diabetes and endocrinology' 'lack of access to pituitary patients and MDTs' 'no community diabetes in 5 years'

Figure 2. Graph to show breakdown of satisfaction ratings for specific aspects of D&E clinical experience





differences in perceptions regarding expectations, regional variations in experience, differences in stage of training and biases from trainees which may or may not be justified. These issues have hopefully been overcome by rolling out the survey at the largest D&E



annual training meeting and achieving a relatively high response rate (20%) and by also publishing the questionnaire online for remote access for those unable to attend.

The balance of registrar workload appears skewed towards GIM, and this is frequently a source of great frustration and disappointment to trainees who are keen to pursue experience in D&E. When compared with other medical specialities such as respiratory medicine or gastroenterology, it would appear that diabetes trainees have an excessive proportion of GIM activity. Whilst recognising the importance of GIM as a specialty, and the benefits of a dual CCT, shortfall in dedicated time to pure specialty training has been highlighted. Consequently, a proportion of D&E trainees feel inadequately prepared or experienced for specialist consultant posts following CCT.

Previous studies have documented a lack of confidence amongst junior doctors in the management of diabetes relative to the management of other common medical conditions.¹² As these junior doctors progress through the specialist training programme, a clear emphasis has to be placedon the acquisition of specialist skills in D&E. Our survey suggests that the pressures of service commitments in GIM adversely affect the opportunities and ability to acquire these vital skills.

It is difficult to attempt to suggest that the achievement of certain curriculum matched competencies, knowledge standards and quantities of clinical attendances translate into a well rounded new consultant by the time of CCT.¹³ Few would disagree that there is a substantial variation in the quality of trainees and new consultants and it is clear that the newly qualified consultant should not be viewed as the



Trainees require:

- more dedicated D&E training time
- a reduction in GIM responsibilities
- closer educational supervision and support
- better access to sub-specialty clinical training
- management and leadership experience

'finished product'. From this survey there appear to be some recurrent problems with the delivery and experience of training for D&E registrars.

Recommendations

Trainees should be supported in four main ways;

- Provision and awareness of existing training opportunities – from pump clinics to clinical commissioning meetings.
- 2. Release from the burden of GIM in line with the other major medical specialites, for at least one year.
- 3. Closer educational supervision and support, which should include regular clinical case discussion, direct observation and feedback on clinical activities and communication both verbal and written, plus opportunities for supported tutorials.
- 4. Exposure to and experience of management and leadership opportunities to

ensure the service developers of the future are adequately prepared to play an active role.

Conclusions

It could be argued that registrars themselves should be proactive in striving to improve their training.¹ There is a need to recognise when things are sub-optimal and to get up and do something about gaps in training. This survey shows variation in training opportunities and quality across the UK and that the disparity of the GIM burden continues. Ultimately, better training will lead to better doctors and better patient care.¹4,¹5

References

- Price HC, George J, Wilmot EG, Maitland R, Knox R, Atkin M, Choudhary P. Taking training into your own hands. *Clin Med* 2010; 10:349-51.
 - http://dx.doi.org/10.7861/clinmedicine.10-4-349
- Grant P, Cheer K, Herring R, et al. CSII experience among higher specialist trainees: a Young Diabetologists and Endocrinologists Forum survey. Practical Diabetes 2013;30:332-5. http://dx.doi.org/10.1002/pdi.1806
- 3. Mason NC, Chaudhuri E, Newbery N, Goddard AF. Training in general medicine are juniors getting enough experience? *Clin Med* 2013;**13**:434-9.
- http://dx.doi.org/10.7861/clinmedicine.13-5-434
 4. Future Hospital Commission. Future hospital.
 Caring for medical patients. Royal College of Physicians, London. September 2013. https://www.rcplondon.ac.uk/sites/default/files/future-hospital-commission-report_0.pdf (accessed August 2014)
- Shape of training. Ensuring the future of excellent patient care. Final report. October 2013. http://www.shapeoftraining.co.uk/static/documents/content/Shape_of_training_FI NAL_Report.pdf_53977887.pdf (accessed August 2014)
- Kar P, Higgins K, Atkin M, Richardson T. The Young Diabetologists Forum (Diabetes UK)/ABCD trainee survey. *Pract Diab Int* 2008;25:323-7.
 - http://dx.doi.org/10.1002/pdi.1296
- Ahluwalia R, Atkin M, Gallen I, et al. Diabetes and endocrinology specialist registrars teaching programmes in the United Kingdomtime to set some quality standards? Results from the Association of British Clinical Diabetologists--Young Diabetologists Forum survey. Diabet Med 2012;29:553-4. http://dx.doi.org/10.1111/j.1464-5491.2011.03501.x
- 8. www.youngdiabetologists.org/ (accessed June 2014) state article/search title to be used.
- Said J, Glendenning J, Leaning D, et al. Oncology Registrar's Forum. 2013 Trainee Survey Report. https://www.rcr.ac.uk/docs/ oncology/pdf/ORF_2013_Survey_Report_FIN AL.pdf (accessed April 2014).
- 10. Herring R. Management of raised glucose, a clinical decision tool to reduce length of stay

- of patients with hyperglycaemia. *Diabet Med* 2013;**30**:81-7.
- http://dx.doi.org/10.1111/dme.12006
- Grant, P. Mustafa, O. Malik, R. Setting up an insulin safety group: a practical approach. Practical Diabetes 2012;29:160-2. http://dx.doi.org/10.1002/pdi.1682
- George JT, Warriner D, McGrane DJ, et al. TOPDOC Diabetes Study Team. Lack of confidence among trainee doctors in the management of diabetes: the Trainees Own Perception of Delivery of Care (TOPDOC) Di-
- abetes Study. *QJM* 2011;**104**:761-6. http://dx.doi.org/10.1093/gjmed/hcr046
- Cheer K, George JT, Grant P, et al. One-third of doctors completing specialist training in diabetes fail to secure a substantive consultant post: young Diabetologists' Forum Survey 2010. Clin Med 2012; 12:244-7. http://dx.doi.org/10.7861/clinmedicine.12-3-244
- Grant P, Nwokolo M. Essential and desirable: making the most of your postgraduate training. Clin Med 2014;14:145-8.
 - http://dx.doi.org/10.7861/clinmedicine.14-2-145
- 15. Kleinman K. Catastrophe and care-giving: the failure of medicine as an art. *The Lancet*, 2008;**371**:22-3.

http://dx.doi.org/10.1016/S0140-6736(08)60057-4

Dr Paul Grant E-mail: drpaul.grant@doctors.org.uk

http://dx.doi.org/10.15277/bjdvd.2014.028 Br J Diabetes Vasc Dis 2014;**14**:120-23



RECRUITING NOW



Do you have patients with type 2 diabetes and obesity needing better control, despite previous liraglutide use?

REVISE-Diabesity is a randomised controlled trial which offers the real chance of improved glycaemic control and reduced weight to enrolled participants, who will be randomised to: 1) Liraglutide 1.8mg, 2) Endobarrier, 3) Endobarrier + liraglutide

Endobarrier is an endosopically inserted duodenal-jejunal bypass liner which early studies suggest might lead to considerable weight loss and improved glycaemic control



To refer patients (Glasgow/ Birmingham / London) please contact:

Dr Piya Sen Gupta, ABCD Research Fellow

Email: revise.diabesity@nhs.net Mobile: 07866319487



Please see the study website (includes selection criteria): http://www.diabetologists-abcd.org.uk/research/endobarrier_study.htm ISRCTN00151053, NCT02055014